arXiv:1211.0920v2 [math.SG] 1 Jan 2013

CERTAIN CIRCLE ACTIONS ON KÄHLER MANIFOLDS

HUI LI

Abstract. Let the circle act holomorphically on a $2n$-dimensional compact Kähler manifold M with moment map $\phi\colon M \to \mathbb{R}$. Assume the critical set of ϕ consists of 3 connected components, the extrema being isolated points. We show that M is equivariantly biholomorphic to $\mathbb{CP}^n$, where $n \geq 2$, or to $\widetilde{G}_2(\mathbb{R}^{n+2})$, the Grassmannian of oriented 2-planes in $\mathbb{R}^{n+2}$, where $n \geq 3$, with a standard circle action; we also show that M is equivariantly symplectomorphic to $\mathbb{CP}^n$, where $n \geq 2$, or to $\widetilde{G}_2(\mathbb{R}^{n+2})$, where $n \geq 3$, with a standard circle action.

1. Introduction

Let (M, ω) be a connected compact symplectic manifold. Assume that M admits a nontrivial[1] Hamiltonian circle action with moment map $\phi\colon M \to \mathbb{R}$. We call M a Hamiltonian S^1-manifold. The moment map $\phi\colon M \to \mathbb{R}$ is a Morse-Bott function whose critical set is exactly the fixed point set M^{S^1} of the circle action.

Let (M, ω) be a compact Hamiltonian S^1-manifold whose even Betti numbers are minimal, i.e., $b_{2i}(M) = 1$ for all $0 \leq 2i \leq \dim(M)$. Some recent work studied certain cases of such Hamiltonian S^1-manifolds. The papers [13, 11] studied the case when $\dim(M) = 6$, and [10, 9] studied the case when M is of any dimension and M^{S^1} consists of two connected components. These papers showed that such Hamiltonian S^1-manifolds have certain geometrical and topological invariants the same as those of the complex projective spaces, the Grassmannian of oriented two planes in some $\mathbb{R}^N$, and some other relatively well known Kähler manifolds. In particular, for two important and interesting cases when $\dim(M) = 6$ and when the fixed points are isolated, [11] determined the equivariant symplectomorphism type of the manifolds. While for the case when M is $2n$-dimensional and M^{S^1} consists of two connected components, one is isolated, by a theorem of Delzant [5], M is equivariantly symplectomorphic to $\mathbb{CP}^n$ with a standard circle action.

In the above case of Delzant's theorem, the other fixed component is necessarily of codimension two in M. The recent paper [7] studied a more

---

Key words and phrases. Kähler manifold, Hamiltonian circle action, biholomorphism, symplectomorphism, equivariant cohomology, equivariant 1st Chern class and Euler class, prequantum line bundle.

2000 classification. 53D05, 53D20, 55N25, 57R20, 32H02, 14J70, 53D50.

[1] We always assume the action is nontrivial unless otherwise stated.

general case: assume (M, ω) is a compact Hamiltonian S^1-manifold such that M^{S^1} consists of two connected components, one of which has codimension two. The authors used certain invariants to classify the equivariant symplectomorphism type of M.

Let (M, ω) be a compact Hamiltonian S^1-manifold. In [10, Section 4], it was shown that if $b_{2i}(M) = 1$ for all $0 \leq 2i \leq \dim(M)$, then the fixed components of the circle action satisfy

$$\sum_{F \subset M^{S^1}} (\dim(F) + 2) = \dim(M) + 2, \tag{1.1}$$

where the sum is over all the fixed set components. While the converse is false, as shown by Example 1.3 below, it is anticipated that when (1.1) holds, the cohomology groups of M are relatively "simple".

In this paper, we consider a compact Kähler manifold M which admits a holomorphic Hamiltonian circle action with moment map $\phi \colon M \to \mathbb{R}$. Assume that M^{S^1} consists of 3 connected components, X, Y, Z, where $\phi(X) < \phi(Y) < \phi(Z)$ and X and Z are isolated points. Using Morse theory (see Lemma 2.4), we can see that X, Y, Z satisfy the equality (1.1). Hence this is a special case of a Hamiltonian S^1-manifold with fixed point set satisfying (1.1). We have two families of such examples as follows.

Example 1.2. Consider $\mathbb{CP}^n$, where $n \geq 2$. As a coadjoint orbit of $SU(n+1)$, it admits a Kähler structure. Consider the effective $S^1 \subset SU(n+1)$ action on $\mathbb{CP}^n$ given by

$$\lambda \cdot [z_0, z_1, \cdots, z_n] = [\lambda^{-l} z_0, \lambda^{l'} z_1, z_2, \cdots, z_n],$$

where $l, l' \in \mathbb{N}$, and $\gcd(l, l') = 1$. The fixed point set $(\mathbb{CP}^n)^{S^1}$ consists of 3 connected components:

$$X = [z_0, 0, \cdots, 0],\ Y = [0, 0, z_2, \cdots, z_n],\ \text{and } Z = [0, z_1, 0, \cdots, 0],$$

where X and Z are isolated points. The action is holomorphic and is Hamiltonian. The weights of the circle action on the normal bundles of X, Y and Z are respectively

$$\big(l, \cdots, l, l + l'\big),\ (-l, l'),\ \big(-l', \cdots, -l', -(l + l')\big).$$

Example 1.3. Consider $\widetilde{G}_2(\mathbb{R}^{n+2})$, the Grassmannian of oriented 2-planes in $\mathbb{R}^{n+2}$, where $n \geq 3$. We have $\dim\big(\widetilde{G}_2(\mathbb{R}^{n+2})\big) = 2n$. As a coadjoint orbit of $SO(n+2)$, it has a Kähler structure. Consider the $S^1 \subset SO(n+2)$ action on $\widetilde{G}_2(\mathbb{R}^{n+2})$ induced by the action on $\mathbb{R}^{n+2} = \mathbb{C} \times \mathbb{R}^n$ given by

$$\lambda \cdot (z, x_1, \dots, x_n) = (\lambda \cdot z, x_1, \dots, x_n)\,.$$

The fixed point set $\big(\widetilde{G}_2(\mathbb{R}^{n+2})\big)^{S^1}$ consists of 3 connected components:

$$X = \mathbb{P}\,(z, 0, \cdots, 0) = \text{pt},\ Y = \widetilde{G}_2\big(0 \times \mathbb{R}^n\big),\ \text{and } Z = \mathbb{P}\,(z, 0, \cdots, 0) = \text{pt},$$

where X and Z correspond to the two orientations on the real 2-plane $(z, 0, \cdots, 0)$. The S^1 action is holomorphic and Hamiltonian. The weights of the circle action on the normal bundles of X, Y and Z are respectively

$$(1, \cdots, 1), \ (-1, 1), \ (-1, \cdots, -1).$$

Our results in Theorems 1.4 and 1.5 equivariantly identify the considered manifold with one of these examples in the complex and symplectic categories.

Theorem 1.4. *Let the circle act holomorphically on a compact* $2n$*-dimensional Kähler manifold* (M, ω, J) *with moment map* $\phi\colon M \to \mathbb{R}$. *Assume the critical set of* ϕ *consists of* 3 *connected components, the extrema being isolated points. Then* M *is* S^1*-equivariantly biholomorphic to* $\mathbb{CP}^n$, *where* $n \geq 2$, *or to* $\widetilde{G}_2(\mathbb{R}^{n+2})$, *where* $n \geq 3$, *with a standard circle action.*

Theorem 1.5. *Let the circle act holomorphically on a compact* $2n$*-dimensional Kähler manifold* (M, ω, J) *with moment map* $\phi\colon M \to \mathbb{R}$. *Assume the critical set of* ϕ *consists of* 3 *connected components, the extrema being isolated points. Then* M *is* S^1*-equivariantly symplectomorphic to* $\mathbb{CP}^n$, *where* $n \geq 2$, *or to* $\widetilde{G}_2(\mathbb{R}^{n+2})$, *where* $n \geq 3$, *with a standard circle action.*

The work of this paper provides a new idea to study Hamiltonian S^1-manifolds which are Kähler and which have fixed point set satisfying (1.1). We hope to be able to use the method to treat more general cases of the fixed point set when the Hamiltonian S^1-manifold is Kähler.

In this work, we mainly use techniques from symplectic geometry, where one main technique is equivariant cohomology and Morse theory; we also use results from algebraic geometry, and use the set up of geometric quantization.

2. Some preliminary results

In this section, we state and prove some preliminary results for symplectic Hamiltonian S^1-manifolds — for the general case and for the case when the Hamiltonian function has 3 critical components, with isolated extrema.

Let (M, ω) be a Hamiltonian S^1-manifold with moment map $\phi\colon M \to \mathbb{R}$, and let F be a fixed component of the S^1 action. Let us set up the following notations.

- $2\lambda_F$: the Morse index of F as a critical set of the Morse-Bott function ϕ;
- $2\lambda_F^+$: the Morse coindex of F for ϕ, or the Morse index of F for $-\phi$;
- Γ_F: the sum of the weights of the S^1 action on the normal bundle N_F to F;
- Λ_F: the product of the weights of the S^1 action on the normal bundle N_F to F;
- Λ_F^-: the product of the weights of the S^1 action on the negative normal bundle N_F^- to F;

- Λ_F^+: the product of the weights of the S^1 action on the positive normal bundle N_F^+ to F.

For an S^1-manifold M, let $H_{S^1}^*(M;R) = H^*(S^\infty \times_{S^1} M; R)$ be the S^1-equivariant cohomology of the manifold M in R coefficient, where R is a coefficient ring. Let $t \in H_{S^1}^2(\mathrm{pt};\mathbb{Z}) = H^2(\mathbb{CP}^\infty;\mathbb{Z})$ be a generator.

2.1. **Some preliminary results on Hamiltonian S^1-manifolds.**

First, we have the following "equivariant extension" of the cohomology class represented by the symplectic form.

Lemma 2.1. [10, Lemma 2.7] *Let the circle act on a connected compact symplectic manifold (M,ω) with moment map $\phi\colon M \to \mathbb{R}$. Let F be a fixed component. Then there exists $\widetilde{u} \in H_{S^1}^2(M;\mathbb{R})$ so that*

$$\widetilde{u}|_{F'} = [\omega|_{F'}] + t\left(\phi(F) - \phi(F')\right)$$

for any fixed component F'. If $[\omega]$ is an integral class, then $\widetilde{u}$ is an integral class.

Now, for a Hamiltonian S^1-manifold M with $H^2(M;\mathbb{R}) = \mathbb{R}$, we use the above lemma to express the 1st Chern class $c_1(M)$ of M in terms of data related to any two fixed components of the action.

Lemma 2.2. *Let the circle act on a connected compact symplectic manifold (M,ω) with moment map $\phi\colon M \to \mathbb{R}$. Assume $H^2(M;\mathbb{R}) = \mathbb{R}$. Then*

$$c_1(M) = \frac{\Gamma_F - \Gamma_{F'}}{\phi(F') - \phi(F)}[\omega],$$

where F and F' are any two fixed components.

Proof. Since $H^2(M;\mathbb{R}) = \mathbb{R}$, $[\omega] \in H^2(M;\mathbb{R})$ is the generator. So t and the $\widetilde{u}$ in Lemma 2.1 are the generators of $H_{S^1}^2(M;\mathbb{R})$. We can write the equivariant first Chern Class $c_1^{S^1}(M)$ of M as

$$c_1^{S^1}(M) = at + b\widetilde{u}, \quad \text{where} \quad a, b \in \mathbb{R}.$$

Then

$$c_1^{S^1}(M)|_f = \Gamma_F t = at, \quad \text{where } f \in F \text{ is a point,} \quad \text{and}$$
$$c_1^{S^1}(M)|_{f'} = \Gamma_{F'} t = at + bt\left(\phi(F) - \phi(F')\right), \quad \text{where } f' \in F' \text{ is a point.}$$

So

$$a = \Gamma_F, \quad b = (\Gamma_{F'} - \Gamma_F)/\left(\phi(F) - \phi(F')\right).$$

Hence

$$c_1^{S^1}(M) = \Gamma_F t + \frac{\Gamma_{F'} - \Gamma_F}{\phi(F) - \phi(F')}\widetilde{u}.$$

Taking the restriction map $H_{S^1}^2(M;\mathbb{R}) \to H^2(M;\mathbb{R})$, we get

$$c_1(M) = \frac{\Gamma_{F'} - \Gamma_F}{\phi(F) - \phi(F')}[\omega].$$

□

The following lemma gives an expression of the equivariant Euler class of the negative normal bundle of a fixed component F when its Morse index is the sum $\sum_{\phi(F')<\phi(F)}(\dim(F')+2)$.

Lemma 2.3. [10, Remark 4.12] *Let the circle act on a connected compact symplectic manifold* (M,ω) *with moment map* $\phi\colon M\to\mathbb{R}$. *Let* F *be a fixed component and assume* $\sum_{\phi(F')<\phi(F)}(\dim(F')+2)=2\lambda_F$. *Then*

$$e^{S^1}(N_F^-)=\Lambda_F^-\prod_{\phi(F')<\phi(F)}\left(t+\frac{[\omega|_F]}{\phi(F')-\phi(F)}\right)^{\frac{1}{2}\dim(F')+1},$$

where $e^{S^1}(N_F^-)$ *is the equivariant Euler class of the negative normal bundle* N_F^- *of* F.

2.2. **The case when the Hamiltonian function has 3 critical components, the extrema being isolated.**

First we show that, the index of the non-extremal critical component of ϕ satisfies the condition of Lemma 2.3, and $H^2(M;\mathbb{R})\cong\mathbb{R}$.

Lemma 2.4. *Let the circle act on a compact symplectic manifold* (M,ω) *with moment map* $\phi\colon M\to\mathbb{R}$ *such that its critical set consists of* 3 *connected components* X, Y *and* Z, *where* $\phi(X)<\phi(Y)<\phi(Z)$, *with* X *and* Z *being isolated points. Then* $2\lambda_Y=2\lambda_Y^+=2$, *and* $H^2(M;\mathbb{R})\cong\mathbb{R}$.

Proof. Since ϕ has 3 critical components, we have $\dim(M)>2$. Since Z is isolated, we have $2\lambda_Z=\dim(M)>2$. The fact that ϕ is a perfect Morse-Bott function gives

$$\dim H^i(M)=\sum_{F\subset M^{S^1}}\dim H^{i-2\lambda_F}(F). \tag{2.5}$$

Then since X is isolated, and $H^2(M;\mathbb{R})\neq 0$, (2.5) gives $2\lambda_Y=2$. Similarly, using $-\phi$, we get $2\lambda_Y^+=2$. Now the fact $H^2(M;\mathbb{R})\cong\mathbb{R}$ follows from these facts and (2.5). □

In the case wnen the S^1 action is semifree, we have the following:

Lemma 2.6. *Let the circle act semifreely on a compact symplectic manifold* (M,ω) *with moment map* $\phi\colon M\to\mathbb{R}$ *such that its critical set consists of* 3 *connected components* X, Y *and* Z, *where* $\phi(X)<\phi(Y)<\phi(Z)$, *with* X *and* Z *being isolated points. Then* $\dim(M)\geq 6$ *and* Y *is not an isolated point.*

Proof. In [14], it was shown that a semifree Hamiltonian circle action on a connected compact $2n$-dimensional symplectic manifold with only isolated fixed points has exactly 2^n fixed points. So under our assumption, Y is not isolated. Together with Lemma 2.4 that $2\lambda_Y=2\lambda_Y^+=2$, we get $\dim(M)=\dim(Y)+2\lambda_Y+2\lambda_Y^+\geq 6$. □

Next, for the case stated, we write $c_1(M)$ slightly differently from that in Lemma 2.2 so that we can identify an integral class.

Lemma 2.7. *Let the circle act on a compact symplectic manifold* (M, ω) *with moment map* $\phi\colon M \to \mathbb{R}$ *such that its critical set consists of* 3 *connected components* X*,* Y *and* Z*, where* $\phi(X) < \phi(Y) < \phi(Z)$*, with* X *and* Z *being isolated points. Then*

$$c_1(M) = \frac{\Gamma_X - \Gamma_Y}{\Lambda_Y^-}\left[\Lambda_Y^- \frac{\omega}{\phi(Y)-\phi(X)}\right], \textit{ where } [\Lambda_Y^- \tfrac{\omega}{\phi(Y)-\phi(X)}] \textit{ is an integral class.}$$

Proof. By Lemma 2.1, we can take $\widetilde{u} \in H^2_{S^1}(M;\mathbb{R})$ so that $\widetilde{u}|_Y = [\omega|_Y]$ and $\widetilde{u}|_X = t\big(\phi(Y) - \phi(X)\big)$. Using this together with Lemmas 2.4 and 2.3, we get

$$e^{S^1}\left(N_Y^-\right) = \Lambda_Y^-\left(t + \frac{[\omega|_Y]}{\phi(X)-\phi(Y)}\right) = \Lambda_Y^-\left(t + \frac{\widetilde{u}|_Y}{\phi(X)-\phi(Y)}\right). \tag{2.8}$$

Since M^{S^1} consists of 3 connected components and Z is isolated, $\dim(M) = 2\lambda_Z \geq 4$, so $\deg\big(e^{S^1}\left(N_Y^-\right)\big) = 2 < 2\lambda_Z$. Let $M^- = \{x \in M \mid \phi(x) < \phi(Z)\}$. Consider the long exact sequence for the pair (M, M^-) in equivariant cohomology with $\mathbb{Z}$ coefficients, since $H^2_{S^1}(M, M^-;\mathbb{Z}) = H^{2-2\lambda_Z}_{S^1}(Z;\mathbb{Z}) = 0$ and $H^3_{S^1}(M, M^-;\mathbb{Z}) = H^{3-2\lambda_Z}_{S^1}(Z;\mathbb{Z}) = 0$, the long exact sequence splits into the short exact sequence

$$0 \to H^2_{S^1}(M;\mathbb{Z}) \to H^2_{S^1}(M^-;\mathbb{Z}) \to 0. \tag{2.9}$$

Consider the class $\widetilde{e^{S^1}\left(N_Y^-\right)} = \Lambda_Y^-\left(t + \frac{\widetilde{u}}{\phi(X)-\phi(Y)}\right) \in H^2_{S^1}(M;\mathbb{R})$. By (2.8), its restriction to Y is $e^{S^1}\left(N_Y^-\right)$; clearly, its restriction to X is 0. By (2.9), $\widetilde{e^{S^1}\left(N_Y^-\right)}$ is the unique class on M having this property. So $e^{S^1}\left(N_Y^-\right)$ is integral implies $\widetilde{e^{S^1}\left(N_Y^-\right)}$ is integral. Taking the restriction map $H^*_{S^1}(M;\mathbb{R}) \to H^*(M;\mathbb{R})$ for the class $\widetilde{e^{S^1}\left(N_Y^-\right)}$, we get that $[\Lambda_Y^- \frac{\omega}{\phi(X)-\phi(Y)}]$ is an integral class.

By Lemma 2.4, $H^2(M;\mathbb{R}) = \mathbb{R}$. Now apply Lemma 2.2 using the fixed components X and Y. □

3. The circle action and the first Chern class of the manifold

In this section, assuming we have a compact Hamiltonian S^1-manifold (M, ω) whose moment map has 3 critical components, the extrema being isolated, we will classify the circle action and express $c_1(M)$ in terms of $[\omega]$ (Proposition 3.3).

An *isotropy submanifold* $M^{\mathbb{Z}_k} \subsetneq M$ is a symplectic submanifold which is not fixed by the S^1 action, but is fixed by the $\mathbb{Z}_k$ action for some $k > 1$.

We need the following lemma in the proof of Proposition 3.3.

Lemma 3.1. [13, Lemma 2.6] *Let the circle act on a compact symplectic manifold* (M, ω). *Let* p *and* q *be fixed points which lie on the same component* N *of* $M^{\mathbb{Z}_k}$ *for some* $k > 1$. *Then the weights of the action at* p *and at* q *are equal modulo* k.

The projection $\pi\colon S^\infty \times_{S^1} M \to \mathbb{CP}^\infty$ induces a natural push forward map $\pi_*\colon H^*_{S^1}(M;\mathbb{Q}) \to H^*(\mathbb{CP}^\infty;\mathbb{Q})$, which is given by "integration over the fiber", denoted $\int_M$. We will need the following theorem due to Atiyah-Bott, and Berline-Vergne [1, 3], in the proof of Proposition 3.3.

Theorem 3.2. *Let the circle act on a compact manifold* M. *Fix a class* $\alpha \in H^*_{S^1}(M;\mathbb{Q})$. *Then as elements of* $\mathbb{Q}(t)$,

$$\int_M \alpha = \sum_{F \subset M^{S^1}} \int_F \frac{\alpha|_F}{e^{S^1}(N_F)},$$

where the sum is over all fixed components, and $e^{S^1}(N_F)$ *is the equivariant Euler class of the normal bundle to* F.

Proposition 3.3. *Let the circle act effectively on a compact* $2n$*-dimensional symplectic manifold* (M, ω) *with moment map* $\phi\colon M \to \mathbb{R}$ *such that its critical set consists of* 3 *connected components* X, Y *and* Z, *where* $\phi(X) < \phi(Y) < \phi(Z)$, *with* X *and* Z *being isolated points. Then the weights of the* S^1 *normal representation at* X, Y *and* Z *are respectively*

$$(1, \cdots, 1), (-1, 1), (-1, \cdots, -1),$$

or

$$(l, \cdots, l, l+l'), (-l, l'), \left(-l', \cdots, -l', -(l+l')\right), \text{ where } l, l' \in \mathbb{N}, \text{ and } \gcd(l, l') = 1.$$

Moreover, if $[\omega]$ *is a primitive integral class, then*

$$c_1(M) = n[\omega] \quad \text{in the first case, and}$$
$$c_1(M) = (n+1)[\omega] \quad \text{in the second case.}$$

Proof. By Lemma 2.4, $2\lambda_Y = 2\lambda_Y^+ = 2$. We have $2\lambda_Z = 2n \geq 4$.

Case 1. Assume the action is semifree. Then the weights of the S^1 normal representations at X, Y and Z are as in the first case. Then using Lemma 2.7, we get

$$c_1(M) = n\left[\frac{\omega}{\phi(Y) - \phi(X)}\right], \quad \text{where } [\tfrac{\omega}{\phi(Y)-\phi(X)}] \in H^2(M;\mathbb{Z}).$$

If $[\omega]$ is primitive integral, then $\phi(Y) - \phi(X) \in \mathbb{N}$ by Lemma 2.1; so $\left[\frac{\omega}{\phi(Y)-\phi(X)}\right] = [\omega]$ and $c_1(M) = n[\omega]$.

Case 2. Assume the action is effective and is not semifree.

Case (2a). Assume there is an isotropy submanifold M_X^Y between X and Y, which is an $M^{\mathbb{Z}_l}$, where $l > 1$, and there is an isotropy submanifold M_Y^Z between Y and Z, which is an $M^{\mathbb{Z}_{l'}}$, where $l' > 1$. We have $\dim(M_X^Y) = \dim(Y) + 2\lambda_Y = \dim(Y) + 2 + \dim(X)$, and similarly $\dim(M_Y^Z) = \dim(Y) +$

$2+\dim(Z)$; moreover, the weights of the normal representation at Y are $(-l, l')$, where $\gcd(l, l')=1$ since the action is effective. Then the weights of the normal representation at X are $(l, \cdots, l, s)$ for some $s\in\mathbb{N}$. The weights of the normal representation at Z must be $(-l', \cdots, -l', -s)$.

By Lemma 2.4, $H^2(M;\mathbb{R})\cong\mathbb{R}$. By rescaling, we may assume that $[\omega]$ is primitive integral. Since $2<2\lambda_Z$, $[\omega|_{\{m\in M\,|\,\phi(m)<\phi(Z)\}}]$ is primitive integral. Since the manifold $\{m\in M\,|\,\phi(m)<\phi(Z)\}$ deformation retracts to M_X^Y (the retract is given by the gradient flow of $-\phi$), $[\omega|_{M_X^Y}]$ is primitive integral. Similarly, since $2<2\lambda_X^+$, $[\omega|_{M_Y^Z}]$ is primitive integral. Using [10, Proposition 6.1] on M_X^Y and on M_Y^Z, we get

$$\phi(Y)-\phi(X)=l \quad \textit{and} \quad \phi(Z)-\phi(Y)=l'.$$

Then using Lemma 2.2 to compute $c_1(M)$ respectively in terms of X, Y and Y, Z, we get

$$s=l+l' \quad \text{and} \quad c_1(M)=(n+1)[\omega].$$

Case (2b). Assume there is an isotropy submanifold M_X^Y between X and Y, which is an $M^{\mathbb{Z}_l}$, where $l>1$, and there is no isotropy submanifold between Y and Z. Then the weights of the normal representation at Y are $(-l, 1)$, and the weights of the normal representation at X are $(l, \cdots, l, s)$ for some $s\in\mathbb{N}$. Similarly as in Case (2a), assume $[\omega]$ is primitive integral, then

$$\phi(Y)-\phi(X)=l.$$

First assume there is no isotropy submanifold between X and Z. Then $s=1$ and the weights of the normal representation at Z are $(-1, \cdots, -1)$. Using Lemma 2.2 to compute $c_1(M)$ respectively in terms of X, Y and Y, Z, it gives a contradiction. Hence there is an isotropy submanifold between X and Z, and there is only one, which corresponds to the weight s (hence $s>1$). So the weights of the normal representation at Z are $(-1, \cdots, -1, -s)$. Using Lemma 3.1 for the fixed sets X and Z, we have $l+1=as$, where $a>0$. Similarly, using the lemma for the fixed sets X and Y, we get $s=1+bl$, where $b>0$. These together give $a=1$ and $b=1$, so

$$s=l+1.$$

Using Lemma 2.2 to compute $c_1(M)$, we get

$$c_1(M)=(n+1)[\omega] \quad \text{and} \quad \phi(Z)-\phi(Y)=1.$$

For the case when there is an isotropy submanifold between Y and Z and there is no isotropy submanifold between X and Y, we can discuss similarly and we have similar conclusion as above.

Case (2c). Assume there are no isotropy submanifolds between X and Y and between Y and Z. Then the weights of the normal representation at Y are $(-1, 1)$. Since the action is not semifree, there is an isotropy submanifold between X and Z. We will show that there is only one isotropy submanifold between X and Z, which is an $M^{\mathbb{Z}_2}$. If $\dim(M)=4$, using Lemma 3.1, this is clear.

Now assume $\dim(M)>4$. Since X and Z are isolated points, each isotropy submanifold between X and Z is a sphere. If W is the set of weights at X, then the set of weights at Z is $-W$. Then $\Gamma_X=-\Gamma_Z$ and $\Lambda_X=\pm\Lambda_Z$. We also have $\Lambda_Y^-=-1$, $\Lambda_Y^+=1$, and $\Gamma_Y=0$. Assume $[\omega]$ is primitive integral. Then by Lemmas 2.7 and 2.1 (using a similar argument as in **Case 1**),

$$c_1(M)=\Gamma_X\,[\omega], \text{ and } \phi(Y)-\phi(X)=1. \tag{3.4}$$

So $c_1(M)|_Y=\Gamma_X\,[\omega|_Y]$. By symmetry or using a similar argument as the above, $\phi(Z)-\phi(Y)=\phi(Y)-\phi(X)=1$. By Lemmas 2.4 and 2.3, $e^{S^1}(N_Y^-)=-t+\frac{[\omega|_Y]}{\phi(Y)-\phi(X)}=-t+[\omega|_Y]$. Using these lemmas for $-\phi$, we get $e^{S^1}(N_Y^+)=t+\frac{[\omega|_Y]}{\phi(Z)-\phi(Y)}=t+[\omega|_Y]$. First assume $\dim(Y)=2(2k+1)$, where $k\geq 0$. Let $c_1^{S^1}(M)$ be the equivariant first Chern class of M. Then $c_1^{S^1}(M)|_X=\Gamma_X t$, $c_1^{S^1}(M)|_Y=c_1(M)|_Y+t-t=\Gamma_X\,[\omega|_Y]$, $c_1^{S^1}(M)|_Z=\Gamma_Z t$. We have $e^{S^1}(N_X)=\Lambda_X t^n$, $e^{S^1}(N_Y)=e^{S^1}(N_Y^-)\cdot e^{S^1}(N_Y^+)$ and $e^{S^1}(N_Z)=\Lambda_Z t^n$. Using Theorem 3.2 to compute

$$\int_M c_1^{S^1}(M)=0,$$

we get

$$\frac{\Gamma_X}{\Lambda_X}-\Gamma_X a+\frac{\Gamma_Z}{\Lambda_Z}=0, \quad \text{where } a=\textstyle\int_Y[\omega|_Y]^{\frac{\dim(Y)}{2}}=\int_Y[\omega|_Y]^{2k+1}\in\mathbb{Z}.$$

Since Y is symplectic, $a\neq 0$. Since $\dim(M)=2(2k+1)+4=2(2k'+1)$, $\Lambda_X=-\Lambda_Z$. Solving the above equation, we get

$$a=\frac{2}{\Lambda_X}\in\mathbb{Z}.$$

Hence $\Lambda_X=2$ ($\Lambda_X\neq 1$ since the action is not semifree) which implies that there can only be one isotropy submanifold between X and Z, which is fixed by $\mathbb{Z}_2$. Next, assume $\dim(Y)=2(2k)$, where $k>0$. We use Theorem 3.2 to compute

$$\int_M\left(c_1^{S^1}(M)\right)^2=0.$$

A similar argument as the above shows that there is only one isotropy submanifold between X and Z, which is fixed by $\mathbb{Z}_2$. Hence, the weights of the action at X are $(1,\cdots,1,2)$ and the weights of the action at Z are $(-1,\cdots,-1,-2)$. Then (3.4) gives $c_1(M)=(n+1)[\omega]$. □

4. Proof of Theorem 1.4

In this section, we prove Proposition 4.2 below and then prove Theorem 1.4. When M is a Kähler manifold satisfying the assumptions of Theorem 1.4, the first Chern class of the manifold given by Proposition 3.3 has important implication.

Let us first recall the following theorem which was proved in [8].

Theorem 4.1. *Let M be a compact Kähler manifold of complex dimension n. If there exists a positive element $\alpha \in H^{1,1}(M;\mathbb{Z})$ such that $c_1(M) = (n+1)\alpha$, then M is biholomorphic to $\mathbb{CP}^n$; if there exists a positive element $\alpha \in H^{1,1}(M;\mathbb{Z})$ such that $c_1(M) = n\alpha$, then M is biholomorphic to a quadratic hypersurface in $\mathbb{CP}^{n+1}$.*

Proposition 4.2. *Let (M, ω, J) be a compact Kähler manifold of complex dimension n, which admits a holomorphic Hamiltonian circle action. Assume that $[\omega]$ is an integral class. If $c_1(M) = (n+1)[\omega]$, then M is S^1-equivariantly biholomorphic to $\mathbb{CP}^n = \mathbb{P}\left(H^0(M;L)\right)$, and if $c_1(M) = n[\omega]$, then M is S^1-equivariantly biholomorphic to a quadratic hypersurface in $\mathbb{CP}^{n+1} = \mathbb{P}\left(H^0(M;L)\right)$, where L is a holomorphic line bundle over M with first Chern class $[\omega]$ and $H^0(M;L)$ is its space of holomorphic sections.*

The proof of Proposition 4.2 is to incorporate the circle action into the proof of Theorem 4.1. The idea of the proof of Theorem 4.1 is as follows. Since $\alpha \in H^{1,1}(M;\mathbb{Z})$, by a result of Kodaira and Spencer, there is a holomorphic line bundle L over M such that $c_1(L) = \alpha$; and since $c_1(L)$ is positive, L is ample by a theorem of Kodaira. Let $H^0(M;L)$ be the complex vector space of holomorphic sections of L over M. The authors of [8] showed the following fact.

Lemma 4.3. *When $c_1(M) = (n+1)c_1(L)$, $\dim H^0(M;L) = n+1$; and when $c_1(M) = nc_1(L)$, $\dim H^0(M;L) = n+2$.*

Moreover, they showed that $H^0(M;L)$ has no base points, i.e., the elements of $H^0(M;L)$ have no common zeros. Hence, if $\mathbb{P}\left(H^0(M;L)\right)$ is the complex projective space defined as the set of hyperplanes through the origin in the vector space $H^0(M;L)$, then we can define a holomorphic mapping

$$j\colon M \to \mathbb{P}\left(H^0(M;L)\right) \quad \text{by} \tag{4.4}$$

$$j(x) = \{s \in H^0(M;L) \mid s(x) = 0\} \quad \text{for each } x \in M.$$

The authors showed that the map j from M to its image is biholomorphic. In the case when $c_1(M) = (n+1)c_1(L)$, $j(M) = \mathbb{P}\left(H^0(M;L)\right) = \mathbb{CP}^n$; and in the case when $c_1(M) = nc_1(L)$, $j(M)$ is a quadratic hypersurface in $\mathbb{P}\left(H^0(M;L)\right) = \mathbb{CP}^{n+1}$.

Now, we make the argument work equivariantly under the assumptions of Proposition 4.2. By the assumptions of Proposition 4.2,

$$c_1(M) = (n+1)[\omega] \quad \text{or} \quad c_1(M) = n[\omega], \quad \text{where } [\omega] \in H^{1,1}(M;\mathbb{Z}) \text{ is positive.}$$

So we have the ample line bundle L as above with $c_1(L) = [\omega]$ and the results in Theorem 4.1.

We choose a Hermitian structure h on L and a Hermitian connection $\bigtriangledown$ such that its curvature form is ω. Then $(L, h, \bigtriangledown)$ is a (pre)quantum line bundle over M.

Let X_M be the vector field generated by the circle action on M, and let $\widetilde{X_M}$ be the horizontal lift of X_M to TL. Then

$$X_L = \widetilde{X_M} + \phi\,\frac{\partial}{\partial\theta}$$

gives an action of the Lie algebra of S^1 on L, where $\frac{\partial}{\partial\theta}$ is the vector field on L generated by the fiberwise multiplication by $e^{i\theta}$.

Let X, Y, Z be the fixed components of the circle action such that $\phi(X) < \phi(Y) < \phi(Z)$. Since $[\omega]$ is an integral class, Lemma 2.1 implies that

$$\phi(Y) - \phi(X) \in \mathbb{N} \quad \text{and} \quad \phi(Z) - \phi(Y) \in \mathbb{N}. \tag{4.5}$$

Since the moment map is defined up to translation by a constant, (4.5) implies that we may assume the moment map values of the fixed point set components are in the integral lattice of $\mathbb{R}$. By [6, Example 6.10], the above action of the Lie algebra of S^1 on L integrates to an S^1 action on L which is compatible with the S^1 action on M. Then S^1 acts on $H^0(M;L)$ which is given by

$$(\lambda \cdot s)\,(x) = \lambda \cdot s(\lambda^{-1} \cdot x), \quad \text{where } \lambda \in S^1 \text{ and } s \in H^0(M;L); \tag{4.6}$$

or infinitesimally given by

$$X \cdot s = -\nabla_{X_M} s + \sqrt{-1}\phi \cdot s, \quad \text{where } X \in \mathrm{Lie}(S^1) \text{ and } s \in H^0(M;L).$$

The S^1 action on $H^0(M;L)$ induces an S^1 action on $\mathbb{P}\left(H^0(M;L)\right)$.

Lemma 4.7. *Under the assumptions of Proposition 4.2, the holomorphic map j in (4.4) can be made to be S^1-equivariant.*

Proof. The above analysis shows that S^1 acts on $H^0(M;L)$ and on $\mathbb{P}\left(H^0(M;L)\right)$. For any $g \in S^1$, and $s \in H^0(M;L)$, by (4.6), we have

$$(g \cdot s)(x) = g \cdot s(g^{-1} \cdot x).$$

So

$$j(g \cdot x) = \{s \in H^0(M;L) \,|\, s(g \cdot x) = 0\} \quad \text{for each } x \in M, \text{ and}$$

$$\begin{aligned} g{\cdot}j(x) &= \{g{\cdot}s \in H^0(M;L) \,|\, g{\cdot}s(x) = 0\} = \{g{\cdot}s \in H^0(M;L) \,|\, g{\cdot}s(g^{-1}g{\cdot}x) = 0\} \\ &= \{g \cdot s \in H^0(M;L) \,|\, (g \cdot s)(g \cdot x) = 0\} \quad \text{for each } x \in M. \end{aligned}$$

Hence

$$j(g \cdot x) = g \cdot j(x).$$

□

Proposition 4.2 follows from Theorem 4.1 and Lemma 4.7.

Remark 4.8. The embedding j in (4.4) may not be a Kähler embedding. In the case when M is a compact homogeneous Kähler manifold which admits a homogeneous very ample quantum line bundle L, for example, when M is $\mathbb{CP}^n$ or $\widetilde{G}_2(\mathbb{R}^{n+2})$, the embedding is Kähler ([2],[4]).

Proof of Theorem 1.4. By Lemma 2.4, we can rescale the Kähler form ω so that $[\omega]$ is primitive integral.

By quotienting out a finite subgroup of S^1 that acts trivially, we may assume that the S^1 action is effective. Then by Proposition 3.3, $c_1(M) = (n+1)[\omega]$, or $c_1(M) = n[\omega]$ and the action is semifree. In the first case, we have $n \geq 2$ (if $n = 1$, then there can only be 2 fixed components), and in the second case, we have $n \geq 3$ by Lemma 2.6.

By Proposition 4.2, M is equivariantly biholomorphic to $\mathbb{P}\left(H^0(M;L)\right) = \mathbb{CP}^n$ or to a quadratic hypersurface in $\mathbb{P}\left(H^0(M;L)\right) = \mathbb{CP}^{n+1}$ through the equivariant holomorphic map j in (4.4). In particular, the circle action on $j(M)$ has 3 fixed components biholomorphic to those on M.

Since the S^1 action on the complex vector space $H^0(M;L)$ is Hamiltonian, the induced S^1 actions on $\mathbb{P}\left(H^0(M;L)\right)$ and on the invariant complex manifold $j(M)$ are Hamiltonian. In the case when $j(M)$ is a quadratic hypersurface in $\mathbb{CP}^{n+1}$, $j(M)$ can be identified with $\widetilde{G}_2\left(\mathbb{R}^{n+2}\right)$. Now Theorem 1.4 follows. ☐

Remark 4.9. Let (M,ω) be a $2n$-dimensional compact Hamiltonian S^1-manifold, such that the fixed point set M^{S^1} consists of two connected components, X and Y with $\dim(X) + \dim(Y) + 2 = \dim(M)$. [10, Theorem 1] showed that, $c_1(M) = (n+1)[\omega]$, or $c_1(M) = n[\omega]$ with $n \geq 3$ odd, if $[\omega]$ is primitive integral. If we assume M is Kähler and the S^1 action preserves the complex structure, then by Proposition 4.2, M is S^1-equivariantly biholomorphic to $\mathbb{CP}^n$ or to $\widetilde{G}_2(\mathbb{R}^{n+2})$ with $n \geq 3$ odd in the latter, with a standard circle action.

See Examples 1.1 and 1.2 in [10] for the standard circle actions on $\mathbb{CP}^n$ and on $\widetilde{G}_2(\mathbb{R}^{n+2})$ satisfying the required conditions.

5. Proof of Theorem 1.5

In this section, we prove Theorem 1.5. With the help of the Kähler condition and Theorem 1.4, we can prove Theorem 1.5 as follows.

Proof of Theorem 1.5. By Theorem 1.4, there is an S^1-equivariant biholomorphism

$$f\colon (M,\omega,J) \to (M',\omega',J'),$$

where (M',ω',J') represents the Kähler manifold $\mathbb{CP}^n$ with $n \geq 2$ or $\widetilde{G}_2\left(\mathbb{R}^{n+2}\right)$ with $n \geq 3$, with the standard structures. By Lemma 2.4, $H^2(M;\mathbb{R}) = \mathbb{R}$. So by rescaling ω, we may assume that ω and $f^*\omega'$ represent the same cohomology class. Consider the family of forms $\omega_t = (1-t)\omega + tf^*\omega'$, where $t \in [0,1]$. Each ω_t is clearly closed. Each ω_t is also nondegenerate: for any point $x \in M$, suppose $X \in T_xM$ is such that $\omega_t(X, Y) = 0$ for all $Y \in T_xM$. In particular, if we take $Y = JX$, then we get $\omega_t(X, JX) = 0$. Using the facts $\omega(X, JX) \geq 0$, $f_*(JX) = J' f_* X$, and $\omega'(f_*X, J'f_*X) \geq 0$, we get $X = 0$. So ω_t is a family of symplectic forms. It represents the same cohomology class and it is S^1-invariant. By Moser's theorem [12] (Moser's argument without the presence of a group action can be adapted to the case when a compact group acts by choosing invariant objects), there is an S^1-equivariant isotopy Φ_t such that $\Phi_t^*\omega_t = \omega$; in particular, $\Phi_1^* f^* \omega' = \omega$. □

References

[1] M. Atiyah and R. Bott, *The moment map and equivariant cohomology*, Topology, **23** (1984), 1-28.

[2] S. Berceanu and M. Schlichenmaier, *Coherent state embeddings, polar divisors and cauchy formulas*, Journal of Geometry and Physics, **34** (2000), 336-358.

[3] N. Berline and M. Vergne, *Classes caractéristiques équivariantes, formule de localisation en cohomologie équivariante*, C. R. Acad. Sci. Paris, **295** (1982), 539-541.

[4] M. Cahen, S. Gutt and J.H. Rawnsley, *Quantization of Kähler manifolds I, geometric interoretation of Berezin's quantization,* J. of Geo. and Phys., **7** (1990), 45-62.

[5] T. Delzant, *Hamiltoniens périodiques et images convexes de l'application moment*, Bull. Soc. Math. France **116** (1988), 315–339.

[6] V. Ginzburg, V. Guillemin and Y. Karshon, *Moment maps, cobordisms, and Hamiltonian group actions*, American Mathematical Society Mathematical Surveys and Monographs **98**, 2002.

[7] J. C. Hausmann and T. Holm, *Simple Hamiltonian manifolds*, arXiv: 1012.4740v1.

[8] S. Kobayashi and T. Ochiai, *Charaterizations of complex projective spaces and hyperquadrics*, J. Math. Kyoto Univ. **13-1** (1973), 31-47.

[9] H. Li, M. Olbermann and D. Stanley, *One connectivity and finiteness of Hamiltonian S^1-manifolds with minimal fixed sets*, arXiv 1010.2505.

[10] H. Li and S. Tolman, *Hamiltonian circle actions with minimal fixed sets*, Int. J. Math., Vol. **23**, no. 8, 2012.

[11] D. McDuff, *Some 6-dimensional Hamiltonian S^1 manifolds*, J. of Topology, **2** (2009), no. 3, 589–623.

[12] J. Moser, *On the volume elements on a manifold*, Trans. Amer. Math. SOC. **120** (1965), 286-294.

[13] S. Tolman, *On a symplectic generalization of Petrie's conjecture*, Trans. Amer. Math. Soc., **362** (2010), 3963-3996.

[14] S. Tolman and J. Weitsman, *On semifree Hamiltonian circle actions with isolated fixed points*, Topology **39** (2000), no. 2, 299-309.

School of mathematical Sciences, Box 173, Suzhou University, Suzhou, 215006, China.

E-mail address: `hui.li@suda.edu.cn`